

WEYL SPECTRA AND WEYL'S THEOREM

YOUNG MIN HAN AND WOO YOUNG LEE

ABSTRACT. “Weyl’s theorem” for an operator on a Hilbert space is a statement that the complement in the spectrum of the “Weyl spectrum” coincides with the isolated eigenvalues of finite multiplicity. In this paper we consider how Weyl’s theorem survives for polynomials of operators and under quasinilpotent or compact perturbations. First, we show that if T is reduced by each of its finite-dimensional eigenspaces then the weyl spectrum obeys the spectral mapping theorem, and further if T is reduction-isoloid then for every polynomial p , Weyl’s theorem holds for $p(T)$. The results on perturbations are as follows. If T is a “finite-isoloid” operator and if K commutes with T and is either compact or quasinilpotent then Weyl’s theorem is transmitted from T to $T + K$. As a non-commutative perturbation theorem, we also show that if the spectrum of T has no holes and at most finitely many isolated points, and if K is a compact operator then Weyl’s theorem holds for $T + K$ when it holds for T .

Introduction. H. Weyl [22] examined the spectra of all compact perturbations $T + K$ of a hermitian operator T and discovered that $\lambda \in \sigma(T + K)$ for every compact operator K if and only if λ is not an isolated eigenvalue of finite multiplicity in $\sigma(T)$. Today this result is known as Weyl’s theorem, and it has been extended from hermitian operators to hyponormal operators and to Toeplitz operators by L. Coburn [7], to several classes of operators including seminormal operators by S. Berberian [2],[3], and to a few classes of Banach space operators [15],[17]. Weyl’s theorem may fail for even the square of T when it holds for T (see [18, Example 1]). In [14], it was shown that Weyl’s theorem holds for polynomials of hyponormal operators. The first aim of this paper is to extend this result via “Berberian” spectra. On the other hand, Weyl’s theorem is liable to fail under “small” perturbations if “small” is interpreted in the sense of compact or quasinilpotent. Recently Weyl’s theorem under small perturbations has been considered in [11],[12],[13], and [18]. The second aim of this paper is to explore how Weyl’s theorem survives under quasinilpotent or compact perturbations.

Throughout this paper let \mathcal{H} denote an infinite dimensional separable Hilbert space. Let $\mathcal{L}(\mathcal{H})$ denote the algebra of bounded linear operators on \mathcal{H} and let $\mathcal{K}(\mathcal{H})$ denote the closed ideal of compact operators on \mathcal{H} . If $T \in \mathcal{L}(\mathcal{H})$ write $\rho(T)$ for the resolvent set of T ; $\sigma(T)$ for the spectrum of T ; $\pi_0(T)$ for the set of eigenvalues of T ; $\pi_{0f}(T)$ for the eigenvalues of finite multiplicity; $\pi_{0i}(T)$ for the eigenvalues of infinite multiplicity. An operator $T \in \mathcal{L}(\mathcal{H})$ is said to be *Fredholm* if $T^{-1}(0)$ and $T(\mathcal{H})^\perp$ are both finite-dimensional. The *index* of a Fredholm operator $T \in \mathcal{L}(\mathcal{H})$, denoted $\text{ind}(T)$, is given by

$$\text{ind}(T) = \dim T^{-1}(0) - \dim T(\mathcal{H})^\perp (= \dim T^{-1}(0) - \dim T^{*-1}(0)).$$

2000 *Mathematics Subject Classification.* Primary 47A10,47A53,47A55

Key words and phrases. Weyl’s theorem, Weyl spectra, Berberian spectra, perturbations, p -hyponormal operators, Toeplitz operators, Unilateral weighted shifts.

This work was supported by the Brain Korea 21 Project.

Typeset by $\mathcal{A}\mathcal{M}\mathcal{S}$ - $\text{T}\mathcal{E}\mathcal{X}$

An operator $T \in \mathcal{L}(\mathcal{H})$ is said to be *Weyl* if it is Fredholm of index zero and *Browder* if it is Fredholm “of finite ascent and descent”: equivalently ([9, Theorem 7.9.3]) if T is Fredholm and $T - \lambda I$ is invertible for sufficiently small $\lambda \neq 0$ in \mathbb{C} . The essential spectrum $\sigma_e(T)$, the Weyl spectrum $\omega(T)$ and the Browder spectrum $\sigma_b(T)$ of $T \in \mathcal{L}(\mathcal{H})$ are defined by

$$\begin{aligned}\sigma_e(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Fredholm}\}; \\ \omega(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Weyl}\}; \\ \sigma_b(T) &= \{\lambda \in \mathbb{C} : T - \lambda I \text{ is not Browder}\};\end{aligned}$$

then (cf. [9])

$$(0.1) \quad \sigma_e(T) \subseteq \omega(T) \subseteq \sigma_b(T) = \sigma_e(T) \cup \text{acc } \sigma(T) \quad \text{and} \quad \omega(T) \subseteq \eta \sigma_e(T),$$

where we write $\text{acc } K$ and ηK for the *accumulation points* and the *polynomially-convex hull*, respectively, of $K \subseteq \mathbb{C}$. If we write $\text{iso } K = K \setminus \text{acc } K$, and ∂K for the topological boundary of K , and

$$(0.2) \quad \pi_{00}(T) := \{\lambda \in \text{iso } \sigma(T) : 0 < \dim (T - \lambda I)^{-1}(0) < \infty\}$$

for the isolated eigenvalues of finite multiplicity, and ([9])

$$(0.3) \quad p_{00}(T) := \sigma(T) \setminus \sigma_b(T)$$

for the *Riesz points* of $\sigma(T)$, then by the punctured neighborhood theorem, i.e., $\partial \sigma(T) \setminus \sigma_e(T) \subseteq \text{iso } \sigma(T)$ (cf. [9],[10]),

$$(0.4) \quad \text{iso } \sigma(T) \setminus \sigma_e(T) = \text{iso } \sigma(T) \setminus \omega(T) = p_{00}(T) \subseteq \pi_{00}(T).$$

We say that *Weyl’s theorem holds for $T \in \mathcal{L}(\mathcal{H})$* if there is equality

$$(0.5) \quad \sigma(T) \setminus \omega(T) = \pi_{00}(T).$$

If $T \in \mathcal{L}(\mathcal{H})$, write $r(T)$ for the spectral radius of T . It is familiar that $r(T) \leq \|T\|$. An operator T is called *normaloid* if $r(T) = \|T\|$ and *isoloid* if $\text{iso } \sigma(T) \subseteq \pi_0(T)$. An operator T is called *reduction-isoloid* if the restriction of T to any reducing subspace is isoloid. It is well-known ([21, Theorem 2]) that every hyponormal operator is reduction-isoloid.

In Section 1, we prove that Weyl spectra, Browder spectra, and “Berberian” spectra all coincide for operators reduced by each of its finite-dimensional eigenspaces. We also use this result to show that if T is reduced by each of its finite-dimensional eigenspaces then Weyl spectrum obeys the spectral mapping theorem, and further if T is reduction-isoloid then for every polynomial p , Weyl’s theorem holds for $p(T)$.

In Section 2, we show that if T is “finite-isoloid” then Weyl’s theorem is transmitted from T to $T + K$ when K is either compact or quasinilpotent and commutes with T , and that if T is a finite-isoloid operator whose spectrum has no holes and at most finitely many isolated points then Weyl’s theorem is transmitted from T to $T + K$ when K is a compact operator. In addition we give their applications to p -hyponormal operators, Toeplitz operators, and unilateral weighted shifts.

1. Berberian Spectra and Weyl’s Theorem. Suppose that $T \in \mathcal{L}(\mathcal{H})$ is reduced by each of its finite-dimensional eigenspaces. If

$$\mathfrak{M} := \bigvee \{(T - \lambda I)^{-1}(0) : \lambda \in \pi_{0f}(T)\},$$

then \mathfrak{M} reduces T . Let $T_1 := T|_{\mathfrak{M}}$ and $T_2 := T|_{\mathfrak{M}^\perp}$. Then we have ([3, Proposition 4.1]) that

- (i) T_1 is a normal operator with pure point spectrum;
- (ii) $\pi_0(T_1) = \pi_{0f}(T)$;
- (iii) $\sigma(T_1) = \text{cl } \pi_0(T_1)$;
- (iv) $\pi_0(T_2) = \pi_0(T) \setminus \pi_{0f}(T) = \pi_{0i}(T)$.

In this case, Berberian ([3, Definition 5.4]) defined

$$(1.0.1) \quad \tau(T) := \sigma(T_2) \cup \text{acc } \pi_{0f}(T).$$

We shall call $\tau(T)$ the *Berberian spectrum* of T . Berberian used the notation $\tau'(T)$ for the concept of (1.0.1). Berberian has also shown that $\tau(T)$ is a nonempty compact subset of $\sigma(T)$. We can, however, show that Weyl spectra, Browder spectra, and Berberian spectra all coincide for operators reduced by each of its finite-dimensional eigenspaces:

Theorem 1.1. *If $T \in \mathcal{L}(\mathcal{H})$ is reduced by each of its finite-dimensional eigenspaces then*

$$(1.1.1) \quad \tau(T) = \omega(T) = \sigma_b(T).$$

Proof. Let \mathfrak{M} be the closed linear span of the eigenspaces $(T - \lambda I)^{-1}(0)$ ($\lambda \in \pi_{0f}(T)$) and write

$$T_1 := T|_{\mathfrak{M}} \quad \text{and} \quad T_2 := T|_{\mathfrak{M}^\perp}.$$

From the preceding arguments it follows that T_1 is normal, $\pi_0(T_1) = \pi_{0f}(T)$ and $\pi_{0f}(T_2) = \emptyset$. For (1.1.1) it will be shown that

$$(1.1.2) \quad \omega(T) \subseteq \tau(T) \subseteq \sigma_b(T)$$

and

$$(1.1.3) \quad \sigma_b(T) \subseteq \omega(T).$$

For the first inclusion of (1.1.2) suppose $\lambda \in \sigma(T) \setminus \tau(T)$. Then $T_2 - \lambda I$ is invertible and $\lambda \in \text{iso } \pi_0(T_1)$. Since also $\pi_0(T_1) = \pi_{0f}(T_1)$, we have that $\lambda \in \pi_{00}(T_1)$. But since T_1 is normal, it follows that $T_1 - \lambda I$ is Weyl and hence so is $T - \lambda I$. This proves the first inclusion. For the second inclusion of (1.1.2) suppose $\lambda \in \sigma(T) \setminus \sigma_b(T)$. Thus $T - \lambda I$ is Browder but not invertible. Observe that the following equality holds with no other restriction on either R or S :

$$(1.1.4) \quad \sigma_b(R \oplus S) = \sigma_b(R) \cup \sigma_b(S) \quad \text{for each } R \in \mathcal{L}(\mathcal{H}_1) \text{ and } S \in \mathcal{L}(\mathcal{H}_2).$$

Indeed if $\lambda \in \text{iso } \sigma(R \oplus S)$ then λ is either an isolated point of the spectra of direct summands or a resolvent element of direct summands, so that if $R - \lambda I$ and $S - \lambda I$ are Fredholm then by (0.4), λ is either a Riesz point or a resolvent element of direct summands, which implies that $\sigma_b(R) \cup \sigma_b(S) \subseteq \sigma_b(R \oplus S)$, and the reverse inclusion is evident. From this we can see that $T_1 - \lambda I$ and $T_2 - \lambda I$ are both Browder. But since $\pi_{0f}(T_2) = \emptyset$, it follows that $T_2 - \lambda I$ is one-one and hence invertible. Therefore $\lambda \in \pi_{00}(T_1) \setminus \sigma(T_2)$, which implies that $\lambda \notin \tau(T)$. This proves the second inclusion of (1.1.2). For (1.1.3) suppose $\lambda \in \sigma(T) \setminus \omega(T)$ and hence $T - \lambda I$ is Weyl but not invertible. Observe that if \mathcal{H}_1 is a Hilbert space and if an operator $R \in \mathcal{L}(\mathcal{H}_1)$ satisfies the equality $\omega(R) = \sigma_e(R)$, then (cf. [11, Theorem 5])

$$(1.1.5) \quad \omega(R \oplus S) = \omega(R) \cup \omega(S) \quad \text{for each Hilbert space } \mathcal{H}_2 \text{ and } S \in \mathcal{L}(\mathcal{H}_2).$$

Since T_1 is normal, applying the equality (1.1.5) to T_1 in place of R gives that $T_1 - \lambda I$ and $T_2 - \lambda I$ are both Weyl. But since $\pi_{0f}(T_2) = \emptyset$, we must have that $T_2 - \lambda I$ is invertible and therefore $\lambda \in \sigma(T_1) \setminus \omega(T_1)$. Thus from Weyl's theorem for normal operators we can see that $\lambda \in \pi_{00}(T_1)$ and hence $\lambda \in \text{iso } \sigma(T_1) \cap \rho(T_2)$, which by (0.4), implies that $\lambda \notin \sigma_b(T)$. This proves (1.1.3) and completes the proof. \square

As applications of Theorem 1.1 we will give several corollaries below.

Corollary 1.2. *If $T \in \mathcal{L}(\mathcal{H})$ is reduced by each of its finite-dimensional eigenspaces then $\sigma(T) \setminus \omega(T) \subseteq \pi_{00}(T)$.*

Proof. This follows at once from Theorem 1.1. \square

Weyl's theorem is not transmitted to dual operators: for example if $T : \ell_2 \rightarrow \ell_2$ is the unilateral weighted shift defined by

$$(1.2.1) \quad Te_n = \frac{1}{n+1}e_{n+1} \quad (n \geq 0),$$

then $\sigma(T) = \omega(T) = \{0\}$ and $\pi_{00}(T) = \emptyset$, and therefore Weyl's theorem holds for T , but fails for its adjoint T^* . We however have:

Corollary 1.3. *If $T \in \mathcal{L}(\mathcal{H})$ is reduced by each of its finite-dimensional eigenspaces and $\text{iso } \sigma(T) = \emptyset$, then Weyl's theorem holds for T and T^* . In this case, $\sigma(T) = \omega(T)$.*

Proof. If $\text{iso } \sigma(T) = \emptyset$, then it follows from Corollary 1.2 that $\sigma(T) = \omega(T)$, which says that Weyl's theorem holds for T . The assertion that Weyl's theorem holds for T^* follows from noting that $\sigma(T)^* = (\sigma(T))^-$, $\omega(T^*) = (\omega(T))^-$ and $\pi_{00}(T^*) = (\pi_{00}(T))^- = \emptyset$. \square

In Corollary 1.3, the condition “ $\text{iso } \sigma(T) = \emptyset$ ” cannot be replaced by the condition “ $\pi_{00}(T) = \emptyset$ ”: for example consider the operator T defined by (1.2.1).

Corollary 1.4 ([2, Theorem]). *If $T \in \mathcal{L}(\mathcal{H})$ is reduction-isoloid and is reduced by each of its finite-dimensional eigenspaces then Weyl's theorem holds for T .*

Proof. In view of Corollary 1.2, it suffices to show that $\pi_{00}(T) \subseteq \sigma(T) \setminus \omega(T)$. Suppose $\lambda \in \pi_{00}(T)$. Then with the preceding notations, $\lambda \in \pi_{00}(T_1) \cap [\text{iso } \sigma(T_2) \cup \rho(T_2)]$. If $\lambda \in \text{iso } \sigma(T_2)$, then since by assumption T_2 is isoloid we have that $\lambda \in \pi_0(T_2)$ and hence $\lambda \in \pi_{0f}(T_2)$. But since $\pi_{0f}(T_2) = \emptyset$, we should have that $\lambda \notin \text{iso } \sigma(T_2)$. Thus $\lambda \in \pi_{00}(T_1) \cap \rho(T_2)$. Since T_1 is normal it follows that $T_1 - \lambda I$ is Weyl and so is $T - \lambda I$; therefore $\lambda \in \sigma(T) \setminus \omega(T)$. \square

If the condition “reduction-isoloid” is replaced by “isoloid” then Corollary 1.4 may fail (see Examples (1) of [2]).

Corollary 1.5. *If $T \in \mathcal{L}(\mathcal{H})$ is reduced by each of its finite-dimensional eigenspaces then*

$$(1.5.1) \quad p(\omega(T)) = \omega(p(T)) \quad \text{for every polynomial } p.$$

Further if T is reduction-isoloid then for every polynomial p , Weyl's theorem holds for $p(T)$.

Proof. We first claim that if T is reduced by each of its finite-dimensional eigenspaces then so is $p(T)$ for any polynomial p : indeed if we write $T = T_1 \oplus T_2$ as in the proof of Theorem 1.1, then $p(T) = p(T_1) \oplus p(T_2)$ shows that $p(T_1)$ is normal and $\pi_{0f}(p(T_2)) \subseteq p(\pi_{0f}(T_2)) = p(\emptyset) = \emptyset$, which implies that $p(T)$ is reduced by each of its finite-dimensional eigenspaces because any normal operator is reduced by each of its finite-dimensional eigenspaces. Therefore the first assertion follows from Theorem 1.1 together with the fact that the Browder spectrum obeys the spectral mapping theorem. The second assertion follows from Theorem 1 of [18] and Corollary 1.4. \square

An operator $T \in \mathcal{L}(\mathcal{H})$ is said to be *p-hyponormal* if $(T^*T)^p - (TT^*)^p \geq 0$ (cf. [1],[4]). If $p = 1$, T is hyponormal and if $p = \frac{1}{2}$, T is semi-hyponormal. In [14, Theorem 2], it was shown that if T is hyponormal then for every polynomial p , Weyl's theorem holds for $p(T)$. We can prove more:

Corollary 1.6. *If $T \in \mathcal{L}(\mathcal{H})$ is p -hyponormal then for every polynomial p , Weyl's theorem holds for $p(T)$.*

Proof. This follows from Corollary 1.5 and the fact that every p -hyponormal operator is isoloid ([6, Theorem 1]) and is reduced by each of its eigenspaces ([5, Theorem 4]). \square

L. Coburn [7, Corollary 3.2] has shown that if $T \in \mathcal{L}(\mathcal{H})$ is hyponormal and $\pi_{00}(T) = \emptyset$, then T is *extremally noncompact*, in the sense that

$$\|T\| = \|\pi(T)\|,$$

where π is the canonical map of $\mathcal{L}(\mathcal{H})$ onto the Calkin algebra $\mathcal{L}(\mathcal{H})/\mathcal{K}(\mathcal{H})$. His proof relies upon the fact that Weyl's theorem holds for hyponormal operators, and hence $\sigma(T) = \omega(T)$ since $\pi_{00}(T) = \emptyset$. Now we can strengthen the Coburn's argument slightly:

Corollary 1.7. *If $T \in \mathcal{L}(\mathcal{H})$ is normaloid and $\pi_{00}(T) = \emptyset$, then T is extremally noncompact.*

Proof. Since $\sigma(T) \subseteq \eta\omega(T) \cup p_{00}(T)$ for any $T \in \mathcal{L}(\mathcal{H})$, we have that $\eta\sigma(T) \setminus \eta\omega(T) \subseteq \pi_{00}(T)$. Thus by our assumption, $\eta\sigma(T) = \eta\omega(T)$. Therefore we can argue that for each compact operator $K \in \mathcal{L}(\mathcal{H})$,

$$\|T\| = r(T) = r_\omega(T) = r_\omega(T + K) \leq r(T + K) \leq \|T + K\|,$$

where $r_\omega(T)$ denotes the ‘‘Weyl spectral radius’’. This completes the proof. \square

Note that if $T \in \mathcal{L}(\mathcal{H})$ is normaloid and $\pi_{00}(T) = \emptyset$, then Weyl's theorem may fail for T ; for example take $\mathcal{H} = \ell_2 \oplus \ell_2$ and $T = U \oplus U^*$, where U is the unilateral shift.

2. Weyl's Theorem Under Small Perturbations. In general Weyl's theorem for $T \in \mathcal{L}(\mathcal{H})$ is not sufficient for Weyl's theorem for $T + K$ with compact (even finite rank) $K \in \mathcal{L}(\mathcal{H})$ commuting with T (cf. [13, Example 2.3]). But if $T \in \mathcal{L}(\mathcal{H})$ is isoloid then Weyl's theorem is transmitted from T to $T + K$ for commuting finite rank operators $K \in \mathcal{L}(\mathcal{H})$ (cf. [13, Theorem 2.2]). But this may fail if ‘‘finite rank’’ is replaced by ‘‘compact’’. In fact Weyl's theorem may fail even if K is both compact and quasinilpotent: for example, take $T = 0$ and K the operator on ℓ_2 defined by $K(x_1, x_2, \dots) = (\frac{x_2}{2}, \frac{x_3}{3}, \frac{x_4}{4}, \dots)$. We will however show that if ‘‘isoloid’’ condition is strengthened slightly then Weyl's theorem is transmitted from T to $T + K$ if K is either a compact or a quasinilpotent operator commuting with T . We begin with:

Proposition 2.1. *If $K \in \mathcal{L}(\mathcal{H})$ is a compact operator commuting with $T \in \mathcal{L}(\mathcal{H})$ then*

$$\pi_{00}(T + K) \subseteq \text{iso}\sigma(T) \cup \rho(T).$$

Proof. Suppose $\lambda \in \pi_{00}(T + K)$. Assume to the contrary that $\lambda \in \text{acc}\sigma(T)$. Observe that $\sigma(T) = \sigma_b(T) \cup p_{00}(T)$ for every $T \in \mathcal{L}(\mathcal{H})$. Since the Browder spectrum is invariant under the commuting compact perturbations ([9, Theorem 7.7.5]), it follows that the difference between $\sigma(T)$ and $\sigma(T + K)$ consists of the difference between $p_{00}(T)$ and $p_{00}(T + K)$. Since by our assumption, $\lambda \in \text{iso}\sigma(T + K) \cap \text{acc}\sigma(T)$, we can find a sequence $\{\lambda_n\}$ of distinct numbers in $p_{00}(T) \setminus p_{00}(T + K)$ satisfying

- (i) $\lim_n \lambda_n = \lambda$;
- (ii) $\sigma := \{\lambda_n\}_{n=1}^\infty \cup \{\lambda\}$ is an isolated part of $\sigma(T)$.

If \mathcal{N} is a neighborhood of σ which contains no other points of $\sigma(T)$, then by using the spectral projection $P = \frac{1}{2\pi i} \int_{\partial\mathcal{N}} (\lambda I - T)^{-1} d\lambda$ corresponding to σ , we can write T as

$$T = \begin{pmatrix} T_1 & 0 \\ 0 & T_2 \end{pmatrix},$$

where $\sigma(T_1) = \sigma$ and $\sigma(T_2) = \sigma(T) \setminus \sigma$. Since $TK = KT$ and $\sigma(T_1) \cap \sigma(T_2) = \emptyset$, it follows from a corollary of Rosenblum's Theorem (cf. [19, Corollary 0.14]) that K admits a matrix representation

$$K = \begin{pmatrix} K_1 & 0 \\ 0 & K_2 \end{pmatrix} \quad \text{with } T_i K_i = K_i T_i \ (i = 1, 2).$$

Since $\lambda \in \pi_{00}(T + K)$ we can suppose that $\dim(T + K - \lambda I)^{-1}(0) := m < \infty$ and hence $\dim(T_1 + K_1 - \lambda I)^{-1}(0) \leq m$. Observe that $\sigma_b(T_1 + K_1) = \sigma_b(T_1) = \{\lambda\}$. But since $\lambda \in \text{iso } \sigma(T_1 + K_1)$, it follows that $\sigma(T_1 + K_1) \setminus \{\lambda\}$ consists of finitely many elements which are its Riesz points. Thus $\sigma(T_1 + K_1) = p_{00}(T_1 + K_1) \cup \{\lambda\}$. Write

$$s := \sum_{z \in p_{00}(T_1 + K_1)} \dim(T_1 + K_1 - zI)^{-1}(0).$$

Then evidently $s < \infty$. On the other hand, since $\lambda_n \in p_{00}(T)$ for every $n = 1, 2, \dots$, using the spectral projections corresponding to the set $\{\lambda_j\}$ for $j = 1, \dots, s + m + 1$, we can write T_1 as

$$T_1 = \begin{pmatrix} T_{11} & & & & \\ & T_{12} & & & 0 \\ & & \ddots & & \\ & & & \ddots & \\ 0 & & & & T_{1,s+m+1} \end{pmatrix} \oplus T_3,$$

where $\sigma(T_{1j}) = \{\lambda_j\}$ for $j = 1, \dots, s + m + 1$, and $\sigma(T_3) = \{\lambda_n\}_{n=s+m+2}^\infty \cup \{\lambda\}$. Note that $\sigma_b(T_{1j}) = \emptyset$ for $j = 1, \dots, s + m + 1$. Therefore each T_{1j} ($j = 1, \dots, s + m + 1$) is a finite dimensional operator because $\sigma_b(S) \neq \emptyset$ for every bounded linear operator S on an infinite dimensional Hilbert space. Since $T_1 K_1 = K_1 T_1$ and the λ_j ($j = 1, \dots, s + m + 1$) are mutually distinct, it again follows from [19, Corollary 0.14] that K_1 can be written as the form

$$K_1 = \begin{pmatrix} K_{11} & & & & \\ & K_{12} & & & 0 \\ & & \ddots & & \\ & & & \ddots & \\ 0 & & & & K_{1,s+m+1} \end{pmatrix} \oplus K_3.$$

Observe that $T_{1j} + K_{1j}$ is a finite dimensional operator for every $j = 1, \dots, s + m + 1$ and $\sigma(T_{1j} + K_{1j}) \subseteq \sigma(T_1 + K_1) = p_{00}(T_1 + K_1) \cup \{\lambda\}$ for every $j = 1, \dots, s + m + 1$. But since

$$\sum_{j=1}^{s+m+1} \sum_{z \in \sigma(T_{1j} + K_{1j})} \dim(T_{1j} + K_{1j} - zI)^{-1}(0) \geq s + m + 1,$$

it follows that $\lambda \in \sigma(T_{1j} + K_{1j})$ for at least $(m + 1)$ j 's, which implies that $\dim(T_1 + K_1 - \lambda I)^{-1}(0) \geq m + 1$, a contradiction. This completes the proof. \square

An operator $T \in \mathcal{L}(\mathcal{H})$ will be said to be *finite-isoloid* if $\text{iso } \sigma(T) \subseteq \pi_{0f}(T)$. Evidently finite-isoloid \Rightarrow isoloid. The converse is not true in general: for example, take $T = 0$. In particular if $\sigma(T)$ has no isolated points then T is finite-isoloid. We now have:

Theorem 2.2. *Suppose $T \in \mathcal{L}(\mathcal{H})$ is finite-isoloid. If Weyl's theorem holds for T then Weyl's theorem holds for $T+K$ if $K \in \mathcal{L}(\mathcal{H})$ commutes with T and is either compact or quasinilpotent.*

Proof. First we assume that K is a compact operator commuting with T . Suppose Weyl's theorem holds for T . We first claim that with no restriction on T ,

$$(2.2.1) \quad \sigma(T+K) \setminus \omega(T+K) \subseteq \pi_{00}(T+K).$$

For (2.2.1), it suffices to show that if $\lambda \in \sigma(T+K) \setminus \omega(T+K)$ then $\lambda \in \text{iso } \sigma(T+K)$. Assume to the contrary that $\lambda \in \text{acc } \sigma(T+K)$. Then we have that $\lambda \in \sigma_b(T+K) = \sigma_b(T)$, so that $\lambda \in \sigma_e(T)$ or $\lambda \in \text{acc } \sigma(T)$. Remember that the essential spectrum and the Weyl spectrum are invariant under compact perturbations. Thus if $\lambda \in \sigma_e(T)$ then $\lambda \in \sigma_e(T+K) \subseteq \omega(T+K)$, a contradiction. Therefore we should have that $\lambda \in \text{acc } \sigma(T)$. But since Weyl's theorem holds for T and $\lambda \notin \omega(T+K) = \omega(T)$, it follows that $\lambda \in \pi_{00}(T)$, a contradiction. This proves (2.2.1). For the reverse inclusion suppose $\lambda \in \pi_{00}(T+K)$. Then by Proposition 2.1, either $\lambda \in \text{iso } \sigma(T)$ or $\lambda \in \rho(T)$. If $\lambda \in \rho(T)$ then evidently $T+K - \lambda I$ is Weyl, i.e., $\lambda \notin \omega(T+K)$. If instead $\lambda \in \text{iso } \sigma(T)$ then $\lambda \in \pi_{00}(T)$ whenever T is finite-isoloid. Since Weyl's theorem holds for T , it follows that $\lambda \notin \omega(T)$ and hence $\lambda \notin \omega(T+K)$. Therefore Weyl's theorem holds for $T+K$.

Next we assume that K is a quasinilpotent operator commuting with T . Then it is known ([18, Lemma 2]) that $\varpi(T) = \varpi(T+K)$ with $\varpi = \sigma, \omega$. Suppose Weyl's theorem holds for T . Then

$$\sigma(T+K) \setminus \omega(T+K) = \sigma(T) \setminus \omega(T) = \pi_{00}(T) \subseteq \text{iso } \sigma(T) = \text{iso } \sigma(T+K),$$

which implies that $\sigma(T+K) \setminus \omega(T+K) \subseteq \pi_{00}(T+K)$. Conversely, suppose $\lambda \in \pi_{00}(T+K)$. If T is finite-isoloid then $\lambda \in \text{iso } \sigma(T+K) = \text{iso } \sigma(T) \subseteq \pi_{0f}(T)$. Thus $\lambda \in \pi_{00}(T) = \sigma(T) \setminus \omega(T) = \sigma(T+K) \setminus \omega(T+K)$. This completes the proof. \square

Corollary 2.3. *Suppose $T \in \mathcal{L}(\mathcal{H})$ is p -hyponormal. If T satisfies one of the following:*

- (i) *iso $\sigma(T) = \emptyset$;*
- (ii) *T has finite-dimensional eigenspaces,*

then Weyl's theorem holds for $T+K$ if $K \in \mathcal{L}(\mathcal{H})$ is either compact or quasinilpotent and commutes with T .

Proof. Observe that each of the conditions (i) and (ii) forces p -hyponormal operators to be finite-isoloid. Since Weyl's theorem holds for p -hyponormal operators ([6]), the result follows at once from Theorem 2.2. \square

It is known ([18, Theorem 3]) that Weyl's theorem is transmitted from $T \in \mathcal{L}(\mathcal{H})$ to $T+K$ for commuting nilpotent operators $K \in \mathcal{L}(\mathcal{H})$. This however does not extend to commuting quasinilpotent operators (see the remark above Proposition 2.1). But if K is an *injective* quasinilpotent operator commuting with T then Weyl's theorem is transmitted from T to $T+K$.

Theorem 2.4. *If Weyl's theorem holds for $T \in \mathcal{L}(\mathcal{H})$ then Weyl's theorem holds for $T+K$ if $K \in \mathcal{L}(\mathcal{H})$ is an injective quasinilpotent operator commuting with T .*

Proof. First of all we prove that if there exists an injective quasinilpotent operator commuting with T , then

$$(2.4.1) \quad T \text{ Weyl} \implies T \text{ injective.}$$

To show this suppose K is an injective quasinilpotent operator commuting with T . Assume to the contrary that T is Weyl but not injective. Then there exists a nonzero vector $x \in \mathcal{H}$ such that

$Tx = 0$. Then by the commutativity assumption, $TK^n x = K^n Tx = 0$ for every $n = 0, 1, 2, \dots$, so that $K^n x \in T^{-1}(0)$ for every $n = 0, 1, 2, \dots$. We now claim that $\{K^n x\}_{n=0}^\infty$ is a sequence of linearly independent vectors in \mathcal{H} . To see this suppose $c_0 x + c_1 Kx + \dots + c_n K^n x = 0$. We may then write $c_n (K - \lambda_1 I) \dots (K - \lambda_n I)x = 0$. Since K is an injective quasinilpotent operator it follows that $(K - \lambda_1 I) \dots (K - \lambda_n I)$ is injective. But since $x \neq 0$ we have that $c_n = 0$. By an induction we also have that $c_{n-1} = \dots = c_1 = c_0 = 0$. This shows that $\{K^n x\}_{n=0}^\infty$ is a sequence of linearly independent vectors in \mathcal{H} . From this we can see $T^{-1}(0)$ is infinite-dimensional, which contradicts to the fact that T is Weyl. This proves (2.4.1). From (2.4.1) we can see that if Weyl's theorem holds for T then $\pi_{00}(T) = \emptyset$. We now claim that $\pi_{00}(T + K) = \emptyset$. Indeed if $\lambda \in \pi_{00}(T + K)$, then $0 < \dim(T + K - \lambda I)^{-1}(0) < \infty$, so that there exists a nonzero vector $x \in \mathcal{H}$ such that $(T + K - \lambda I)x = 0$. But since K commutes with $T + K - \lambda I$, the same argument as in the proof of (2.4.1) with $T + K - \lambda I$ in place of T shows that $(T + K - \lambda I)^{-1}(0)$ is infinite-dimensional, a contradiction. Therefore $\pi_{00}(T + K) = \emptyset$ and hence Weyl's theorem holds for $T + K$ because $\varpi(T) = \varpi(T + K)$ with $\varpi = \sigma, \omega$. \square

In Theorem 2.4, “quasinilpotent” cannot be replaced by “compact”. For example consider the following operators on $\ell_2 \oplus \ell_2$:

$$T = \begin{pmatrix} 0 & & & & \\ \frac{1}{2} & & 0 & & \\ & \frac{1}{3} & & & \\ 0 & & \frac{1}{4} & & \\ & & & \ddots & \end{pmatrix} \oplus I \quad \text{and} \quad K = \begin{pmatrix} 1 & & & & \\ -\frac{1}{2} & & 0 & & \\ & -\frac{1}{3} & & & \\ 0 & & -\frac{1}{4} & & \\ & & & \ddots & \end{pmatrix} \oplus Q,$$

where Q is an injective compact quasinilpotent operator on ℓ_2 . Observe that Weyl's theorem holds for T , K is an injective compact operator, and $TK = KT$. But

$$\sigma(T + K) = \{0, 1\} = \omega(T + K) \quad \text{and} \quad \pi_{00}(T + K) = \{1\},$$

which says that Weyl's theorem does not hold for $T + K$.

In the perturbation theory the “commutative” condition looks so rigid. Without the commutativity, the spectrum can however undergo a large change under even rank one perturbations. In spite of it, Weyl's theorem may hold for (non-commutative) compact perturbations of “good” operators. We now give such a perturbation theorem. To do this we need:

Lemma 2.5. *If $N \in \mathcal{L}(\mathcal{H})$ is a quasinilpotent operator commuting with $T \in \mathcal{L}(\mathcal{H})$ modulo compact operators (i.e., $TN - NT \in \mathcal{K}(\mathcal{H})$) then $\sigma_e(T + N) = \sigma_e(T)$ and $\omega(T + N) = \omega(T)$.*

Proof. Straightforward from [18, Lemma 2]. \square

Theorem 2.6. *Suppose $T \in \mathcal{L}(\mathcal{H})$ satisfies the following:*

- (i) T is finite-isoloid;
- (ii) $\sigma(T)$ has no “holes” (bounded components of the complement), i.e., $\sigma(T) = \eta \sigma(T)$;
- (iii) $\sigma(T)$ has at most finitely many isolated points;
- (iv) Weyl's theorem holds for T .

If $K \in \mathcal{L}(\mathcal{H})$ is either compact or quasinilpotent and commutes with T modulo compact operators then Weyl's theorem holds for $T + K$.

Proof. By Lemma 2.5, we have that $\sigma_e(T + K) = \sigma_e(T)$ and $\omega(T + K) = \omega(T)$. Suppose Weyl's theorem holds for T and $\lambda \in \sigma(T + K) \setminus \omega(T + K)$. We now claim that $\lambda \in \text{iso } \sigma(T + K)$. Assume to the contrary that $\lambda \in \text{acc } \sigma(T + K)$. Since $\lambda \notin \omega(T + K) = \omega(T)$, it follows from

the punctured neighborhood theorem that $\lambda \notin \partial\sigma(T + K)$. Also since the set of all Weyl operators forms an open subset of $\mathcal{L}(\mathcal{H})$, we have that $\lambda \in \text{int}(\sigma(T + K) \setminus \omega(T + K))$. Then there exists $\epsilon > 0$ such that $\{\mu \in \mathbb{C} : |\mu - \lambda| < \epsilon\} \subseteq \text{int}(\sigma(T + K) \setminus \omega(T + K))$, and hence $\{\mu \in \mathbb{C} : |\mu - \lambda| < \epsilon\} \cap \omega(T) = \emptyset$. But since

$$\partial\sigma(T + K) \setminus \text{iso}\sigma(T + K) \subseteq \sigma_e(T + K) = \sigma_e(T),$$

it follows from our assumption that

$$\begin{aligned} \{\mu \in \mathbb{C} : |\mu - \lambda| < \epsilon\} &\subseteq \text{int}(\sigma(T + K) \setminus \omega(T + K)) \\ &\subseteq \eta(\partial\sigma(T + K) \setminus \text{iso}\sigma(T + K)) \\ &\subseteq \eta\sigma_e(T) \subseteq \eta\sigma(T) = \sigma(T), \end{aligned}$$

which implies that $\{\mu \in \mathbb{C} : |\mu - \lambda| < \epsilon\} \subseteq \sigma(T) \setminus \omega(T)$. This contradicts to Weyl's theorem for T . Therefore $\lambda \in \text{iso}\sigma(T + K)$ and hence $\sigma(T + K) \setminus \omega(T + K) \subseteq \pi_{00}(T + K)$. For the reverse inclusion suppose $\lambda \in \pi_{00}(T + K)$. Assume to the contrary that $\lambda \in \omega(T + K)$ and hence $\lambda \in \omega(T)$. Then we claim $\lambda \notin \partial\sigma(T)$. Indeed if $\lambda \in \text{iso}\sigma(T)$ then by assumption $\lambda \in \pi_{00}(T)$, which contradicts to Weyl's theorem for T . If instead $\lambda \in \text{acc}\sigma(T) \cap \partial\sigma(T)$ then since $\text{iso}\sigma(T)$ is finite it follows that

$$\lambda \in \text{acc}(\partial\sigma(T)) \subseteq \text{acc}\sigma_e(T) = \text{acc}\sigma_e(T + K),$$

which contradicts to the fact that $\lambda \in \text{iso}\sigma(T + K)$. Therefore $\lambda \notin \partial\sigma(T)$. Also since $\lambda \in \text{iso}\sigma(T + K)$, there exists $\epsilon > 0$ such that

$$\{\mu \in \mathbb{C} : 0 < |\mu - \lambda| < \epsilon\} \subseteq \sigma(T) \cap \rho(T + K),$$

so that $\{\mu \in \mathbb{C} : 0 < |\mu - \lambda| < \epsilon\} \cap \omega(T) = \emptyset$, which contradicts to Weyl's theorem for T . Thus $\lambda \in \sigma(T + K) \setminus \omega(T + K)$ and therefore Weyl's theorem holds for $T + K$. \square

If, in Theorem 2.6, the condition “ $\sigma(T)$ has no holes” is dropped then Theorem 2.6 may fail even though T is normal. For example, if on $\ell_2 \oplus \ell_2$

$$T = \begin{pmatrix} U & I - UU^* \\ 0 & U^* \end{pmatrix} \quad \text{and} \quad K = \begin{pmatrix} 0 & I - UU^* \\ 0 & 0 \end{pmatrix},$$

where U is the unilateral shift on ℓ_2 , then T is unitary (essentially the bilateral shift) with $\sigma(T) = \mathbb{T}$, K is a rank one nilpotent, and Weyl's theorem does not hold for $T - K$.

Also in Theorem 2.6, the condition “ $\text{iso}\sigma(T)$ is finite” is essential in the cases where K is compact. For example, if on ℓ_2

$$T(x_1, x_2, \dots) = (x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots) \quad \text{and} \quad Q(x_1, x_2, \dots) = (\frac{x_2}{2}, \frac{x_3}{3}, \frac{x_4}{4}, \dots),$$

we define $K := -(T + Q)$. Then we have that (i) T is finite-isoloid; (ii) $\sigma(T)$ has no holes; (iii) Weyl's theorem holds for T ; (iv) $\text{iso}\sigma(T)$ is infinite; (v) K is compact because T and Q are both compact; (vi) Weyl's theorem does not hold for $T + K (= -Q)$.

Corollary 2.7. *If $\sigma(T)$ has no holes and at most finitely many isolated points and if K is a compact operator then Weyl's theorem is transmitted from T to $T + K$.*

Proof. Straightforward from Theorem 2.6. □

Corollary 2.7 shows that if Weyl's theorem holds for T whose spectrum has no holes and at most finitely many isolated points then for every compact operator K , the passage from $\sigma(T)$ to $\sigma(T + K)$ is putting at most countably many isolated points outside $\sigma(T)$ which are Riesz points of $\sigma(T + K)$. Here we should note that this holds even if T is quasinilpotent because for every quasinilpotent operator T (more generally, "Riesz operators"), we have

$$\sigma(T + K) \subseteq \eta \sigma_e(T + K) \cup p_{00}(T + K) = \eta \sigma_e(T) \cup p_{00}(T + K) = \{0\} \cup p_{00}(T + K).$$

Corollary 2.7 can easily be applied for Toeplitz operators and unilateral weighted shifts. In the below we give two corollaries on those operators. Let $H^2(\mathbb{T})$ denote the Hardy space of the unit circle $\mathbb{T} = \partial\mathbb{D}$ in the complex plane. Recall ([16]) that given $\varphi \in L^\infty(\mathbb{T})$, the *Toeplitz operator* with symbol φ is the operator T_φ on $H^2(\mathbb{T})$ defined by $T_\varphi f = P(\varphi \cdot f)$, where $f \in H^2(\mathbb{T})$ and P denotes the projection that maps $L^2(\mathbb{T})$ onto $H^2(\mathbb{T})$. For example if $\varphi(z) = z$ then T_φ represents the unilateral shift on ℓ_2 . Write $C(\mathbb{T})$ for the algebra of all continuous complex-valued functions on \mathbb{T} .

Corollary 2.8. *Suppose T_φ is a Toeplitz operator with nonconstant continuous symbols $\varphi \in C(\mathbb{T})$ whose winding number of φ with respect to each hole of $\varphi(\mathbb{T})$ is nonzero. If $K \in \mathcal{L}(H^2)$ is a compact operator then Weyl's theorem holds for $T_\varphi + K$. Hence, in particular, if U is the unilateral shift on ℓ_2 then Weyl's theorem holds for $U + K$ with every compact operator $K \in \mathcal{L}(\ell_2)$.*

Proof. Remember ([7]) that Weyl's theorem holds for every Toeplitz operator and $\sigma(T_\varphi)$ has no isolated points for nonconstant symbols φ . The spectral theory for Toeplitz operators with continuous symbols shows that our assumption implies $\sigma(T_\varphi)$ has no holes (cf. [16]). Therefore the result follows at once from Corollary 2.7. The second assertion is immediate from the first. □

Corollary 2.9. *If T is a unilateral weighted shift with positive weights and is not quasinilpotent, and if $K \in \mathcal{L}(\ell_2)$ is a compact operator then Weyl's theorem holds for $T + K$.*

Proof. If T is a unilateral weighted shift which is not quasinilpotent then $\sigma(T)$ is a non-degenerated disk ([20, Theorem 4]). Moreover since weights are positive it follows that $\pi_0(T) = \emptyset$, and hence $\sigma(T) = \omega(T)$, which implies that Weyl's theorem holds for T . Therefore the result follows at once from Corollary 2.7. □

Corollary 2.9 may fail if T is quasinilpotent: for example consider the operators T and K defined by $T(x_1, x_2, \dots) = (0, x_1, \frac{x_2}{2}, \frac{x_3}{3}, \dots)$ and $K(x_1, x_2, \dots) = (0, -x_1, 0, 0, \dots)$.

Acknowledgement. The authors wish to thank Professor Robin Harte for helpful discussions concerning this work.

REFERENCES

1. A. Aluthge, *On p -hyponormal operators for $0 < p < 1$* , Integral Equations Operator Theory **13** (1990), 307–315.
2. S.K. Berberian, *An extension of Weyl's theorem to a class of not necessarily normal operators*, Michigan Math. J. **16** (1969), 273–279.
3. S.K. Berberian, *The Weyl spectrum of an operator*, Indiana Univ. Math. J. **20** (1970), 529–544.

4. M. Chō, *Spectral properties of p -hyponormal operators*, Glasgow Math. J. **36** (1994), 117–122.
5. M. Chō and T. Huruya, *p -hyponormal operators ($0 < p < \frac{1}{2}$)*, Commentationes Math. **33** (1993), 23–29.
6. M. Chō, M. Itoh and S. Ōshiro, *Weyl's theorem holds for p -hyponormal operators*, Glasgow Math. J. **39** (1997), 217–220.
7. L.A. Coburn, *Weyl's theorem for nonnormal operators*, Michigan Math. J. **13** (1966), 285–288.
8. R.E. Harte, *Fredholm, Weyl and Browder theory*, Proc. Royal Irish Acad. **85A (2)** (1985), 151–176.
9. R.E. Harte, *Invertibility and Singularity for Bounded Linear Operators*, Dekker, New York, 1988.
10. R.E. Harte and W.Y. Lee, *The punctured neighbourhood theorem for incomplete spaces*, J. Operator Theory **30** (1993), 217–226.
11. R.E. Harte and W.Y. Lee, *Another note on Weyl's theorem*, Trans. Amer. Math. Soc. **349** (1997), 2115–2124.
12. W.Y. Lee, *Weyl's theorem for operator matrices*, Integral Equations Operator Theory **32** (1998), 319–331.
13. W.Y. Lee and S.H. Lee, *On Weyl's theorem (II)*, Math. Japo. **43** (1996), 549–553.
14. W.Y. Lee and S.H. Lee, *A spectral mapping theorem for the Weyl spectrum*, Glasgow Math. J. **38** (1996), 61–64.
15. V.I. Istrătescu, *On Weyl's spectrum of an operator. I*, Rev. Roum. Math. Pures Appl. **17** (1972), 1049–1059.
16. N.K. Nikolskii, *Treatise on the Shift Operators*, Springer, New York, 1986.
17. K.K. Oberai, *On the Weyl spectrum*, Illinois J. Math. **18** (1974), 208–212.
18. K.K. Oberai, *On the Weyl spectrum (II)*, Illinois J. Math. **21** (1977), 84–90.
19. H. Radjavi and P. Rosenthal, *Invariant Subspaces*, Springer, New York, 1973.
20. A.L. Shields, *Weighted shift operators and analytic function theory*, Mathematical Surveys **13** (1974), 49–128 (C. Pearcy, ed., *Topics in Operator Theory*, A.M.S. Providence, 1974).
21. J. Stampfli, *Hyponormal operators*, Pacific J. Math. **12** (1962), 1453–1458.
22. H. Weyl, *Über beschränkte quadratische Formen, deren Differenz vollsteig ist*, Rend. Circ. Mat. Palermo **27** (1909), 373–392.

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 440-746, KOREA
E-mail address: ymhan@math.skku.ac.kr

DEPARTMENT OF MATHEMATICS, SUNGKYUNKWAN UNIVERSITY, SUWON 440-746, KOREA
E-mail address: wylee@yurim.skku.ac.kr